

Poisonous Plants Policy

Rationale:

Te Rāwhiti Kindergarten is committed to ensuring tamariki are kept safe from harm and upholding Ministry of Education Licensing Criteria HS12 in which hazards are eliminated, isolated or minimized.

Te Whāriki:

Well-being/Mana Atua: Tamariki experience an environment where they are kept safe from harm.

Procedures:

1. A list of plants poisonous to tamariki from Manaaki Whenua Landcare Research is attached to this policy.
2. The listed plants are not to be bought or planted within the kindergarten gardens.
3. Kindergarten kaiako are aware of the poisonous nature of the listed plants.
4. Kindergarten kaiako supervise any gathering of flowers, fruits or vegetables by the tamariki.
5. Kindergarten kaiako ensure that there is no unsupervised eating of anything whilst tamariki play in the garden and on trips.
6. Tamariki wash hands after playing in the garden.
7. Kindergarten kaiako have current First Aid qualifications in line with Licensing Criteria HS25 (One adult to every 25 tamariki).
8. For any concerns around exposure to poisonous plants, kaiako will ring the National Poisons Centre on 0800 764 766 (0800 POISON).

Authorised:	Shelley W
Date:	15 Dec 2021
Review Date:	15 Dec 2023
Consultation Undertaken:	30 Sept – 17 Oct 2021



Safety in Pre-school Centres

Plants which existing literature suggests should not be grown or tolerated in pre-school centres

INTERNAL POISONS—Harmful if swallowed

- Angel's trumpet (*Brugmansia candida*) – North Island and warmest parts of South Island
- Arums and arum lily (*Arum* species and *Zantedeschia aethiopica*)
- Bittersweet (*Solanum dulcamara*) – mainly South Island and lower North Island
- Castor oil (*Ricinus communis*)
- Death cap and fly agaric fungi (*Agaricus phalloides* and *A. muscaria*)
- Fox glove (*Digitalis purpurea*)
- Hemlock (*Conium maculatum*)
- Jerusalem cherry (*Solanum diflorum* and *S. pseudocapsicum*)
- Laburnum (*Laburnum anagyroides*) – mainly South Island and southern half of North Island
- Lantana (*Lantana camara*) – warmer parts of North Island and northern areas of South Island
- Lily of the valley (*Convallaria majalis*) – mainly South Island
- Monkshood (*Aconitum napellus*) – mainly South Island and colder parts of North Island
- Oleander (*Nerium oleander*)
- Persian lilac or white cedar (*Melia azederach*) – mainly North Island
- Potato (*Solanum tuberosum*) – all green parts
- Privet species (*Ligustrum* species)
- Queen of the night (*Cestrum nocturnum*) – mainly northern North Island
- Spindle tree and Japanese spindle tree (*Fuonymus europaeus* and *F. japonicus*) – latter mainly fruiting in North Island
- Stinking iris (*Iris foetidissima*)
- Tutu (*Coriaria* species) – nearly always *C. arborea*
- Yew (*Taxus baccata*) – although nearly all parts of both sexes are poisonous, only the berry of the female tree, with its poisonous seed, is likely to be eaten

EXTERNAL POISONS—Harmful if touched

- Stinging nettles (*Urtica* species)
- Wax tree or Japanese wax tree (*Rhus succedanea*) – mainly North Is. and northern South Is.

Notes

Certain very poisonous plants are excluded from the above list because of their rarity in New Zealand, e.g. deadly nightshade* (*Atropa belladonna*), bushman's poison (*Acokanthera oppositifolia*), henbane (*Hyoscyamus niger*), and poison ivy (*Rhus radicans*). Also excluded are poisonous plants which, although common, have no parts likely to attract young children; e.g. hellebores (*Helleborus* species), box (*Buxus sempervirens*), and thornapple or datura (*Datura stramonium*), the last having very poisonous seeds enclosed in spiny capsules.

Many plants have spiny or prickly vegetative parts but these are not considered to be reason enough to exclude such plants from pre-school education centres. Most of these plants are either not poisonous or not very poisonous; e.g. roses (*Rosa* species and hybrids), firethorns, (*Pyracantha* species), hawthorn (*Crataegus monogyna*), gooseberry (*Ribes uva-crispa*), and holly (*Ilex aquifolium*).

* True deadly nightshade is extremely rare in New Zealand. Unfortunately, this name is mistakenly applied to black nightshade (*Solanum nigrum*), the latter having somewhat poisonous green parts and harmless berries. The two are easily distinguished: deadly nightshade is a large plant over a metre high when mature, and has large, bell-shaped, brownish-purple flowers followed by large, egg-shaped black berries. Black nightshade is a much smaller plant (about half the height of deadly nightshade when mature), and has small white star-shaped flowers followed by little black berries, similar to black currants.

© Landcare Research New Zealand Ltd 2005. This information may be copied and distributed, provided Landcare Research is acknowledged. Under no circumstances may a charge be made for this information without the permission of Landcare Research.

Disclaimer of Liability for Health-Related Advice

This information is not intended to be comprehensive or to provide medical advice to you. While all care has been taken to ensure the accuracy of the information, no responsibility is accepted for any actions taken in reliance on it. Specific advice on medical issues should be sought from a health professional.

For more information contact: The Plant Herbarium, Landcare Research, P O Box 40, Lincoln 7640,
Tel 03-321 9999, Fax 03-321 9998, Email chr@landcareresearch.co.nz

This sheet is available from the Landcare Research Website: www.landcareresearch.co.nz